

Low-energy electron scattering in a strong Coulomb field

A.I.Milstein, I.S.Terekhov

Budker Institute of Nuclear Physics, 630090 Novosibirsk, Russia

(Dated: February 2, 2008)

Abstract

The analytic expression for the cross section of low-energy electron scattering in a strong Coulomb field is obtained. It is shown that in a wide energy region this cross section differs essentially from that obtained in the first Born approximation.

I. INTRODUCTION

The explicit form of the cross section for electron scattering in a strong Coulomb field at arbitrary energy was obtained many years ago [1]. This form contains the infinite series with respect to the angular momentum. Though various approaches to the summation over the angular momentum were developed in numerous papers, the numerical calculation of the cross section with the use of the results of [1] still rather complicated problem. The detailed review of the papers devoted to the problem under discussion can be found in [2]. In the papers [3, 4, 5] the numerical calculations of the cross section were performed for various scattering angles, the nuclear charge numbers Z , and the kinetic electron energy above 0.023MeV. For $Z = 80$ in the case of backward electron scattering, it was shown that the ratio of the exact cross section and the Rutherford cross section increases from 0.15 to 2.35 when the electron kinetic energy decreases from 1.675MeV to 0.023MeV. Such big deviation from the non-relativistic result stimulate the interest to the investigation of the exact cross section at very small energy. In the present paper we have solved this problem calculating the asymptotic form of the cross section for arbitrary Z and small kinetic electron energy.

II. CROSS SECTION

A simple way to derive the scattering amplitude is the use of the Green function $G(\mathbf{r}_2, \mathbf{r}_1|\varepsilon)$ of the Dirac equation in the external field. The wave function $\psi_{\lambda\mathbf{p}}(\mathbf{r})$ can be obtained with the help of the relation

$$\lim_{r_1 \rightarrow \infty} G(\mathbf{r}_2, \mathbf{r}_1|\varepsilon) = -\frac{\exp(ipr_1)}{4\pi r_1} \sum_{\lambda=1,2} \psi_{\lambda\mathbf{p}}^{(+)}(\mathbf{r}_2) \bar{u}_{\lambda\mathbf{p}}, \quad u_{\lambda\mathbf{p}} = \sqrt{\varepsilon + m} \begin{pmatrix} \phi_\lambda \\ \frac{\boldsymbol{\sigma} \cdot \mathbf{p}}{\varepsilon + m} \phi_\lambda \end{pmatrix}, \quad (1)$$

where $p = \sqrt{\varepsilon^2 - m^2}$, m is the electron mass, $\psi_{\lambda\mathbf{p}}^{(+)}(\mathbf{r})$ denotes a solution of the Dirac equation containing at the infinity a plane wave with the momentum $\mathbf{p} = -p\mathbf{n}_1$ ($\mathbf{n}_{1,2} = \mathbf{r}_{1,2}/r_{1,2}$) and a diverging spherical wave, λ is the helicity, $\hbar = c = 1$. In the Coulomb field the right-hand-side of (1) contains the additional factor $(2pr_1)^{iq}$, where $q = Z\alpha\varepsilon/p$, $\alpha = 1/137$ is the fine-structure constant. In [6] a convenient integral representation was obtained for the electron Green function in the Coulomb field. Using Eqs. (19)-(22) of that paper, we arrive

at the following result:

$$\begin{aligned}
\psi_{\lambda\mathbf{p}}^{(+)}(\mathbf{r}_2) &= \sqrt{\varepsilon + m} \sum_{l=1}^{\infty} \left(\frac{f_1}{\frac{p\boldsymbol{\sigma} \cdot \mathbf{n}_2}{\varepsilon + m} f_2} \right), \\
f_{1,2} &= [(R_1 A + i \frac{mZ\alpha}{p} R_2 B) M_1 \mp i R_2 B M_2] \phi_{\lambda}, \\
A &= l \frac{d}{dx} (P_l(x) + P_{l-1}(x)) \quad , \quad B = \frac{d}{dx} (P_l(x) - P_{l-1}(x)) \quad , \\
x &= \mathbf{n}_1 \cdot \mathbf{n}_2 \quad , \quad R_{1,2} = 1 \mp (\boldsymbol{\sigma} \cdot \mathbf{n}_2)(\boldsymbol{\sigma} \cdot \mathbf{n}_1) \\
M_{1,2} &= i \frac{\exp(ipr_2 - i\pi\nu)}{pr_2} \int_0^{\infty} t^{(\mp 1 - 2iq)} e^{it^2} J_{2\nu}(2t\sqrt{2pr_2}) dt. \tag{2}
\end{aligned}$$

Here $P_l(x)$ is the Legendre polynomial, $J_{2\nu}$ is the Bessel function, $\nu = \sqrt{l^2 - (Z\alpha)^2}$. The integral in the expression for $M_{1,2}$ are expressed via the confluent hypergeometric function. The result (2) is in agreement with the well-known solution of the Dirac equation in the Coulomb field.

When $r_2 \rightarrow \infty$ then the coefficient W_{λ} at the diverging spherical wave

$$\exp[ipr_2 + iq \ln(2pr_2)]/r_2$$

in $\psi_{\lambda\mathbf{p}}^{(+)}(\mathbf{r}_2)$ is determined by the asymptotics of the function M_1 in (2) coming from the region of integration $t \ll 1$. We have

$$\begin{aligned}
W_{\lambda} &= \sqrt{\varepsilon + m} \sum_{l=1}^{\infty} \left(\frac{f}{\frac{p\boldsymbol{\sigma} \cdot \mathbf{n}_2}{\varepsilon + m} f} \right), \\
f &= \frac{ie^{-i\pi\nu}\Gamma(\nu - iq)}{2p\Gamma(\nu + 1 + iq)} [R_1 A + i \frac{mZ\alpha}{p} R_2 B] \phi_{\lambda}. \tag{3}
\end{aligned}$$

As a result we obtain the cross section

$$\begin{aligned}
\frac{d\sigma}{d\Omega} &= \frac{2}{p^2} \left\{ (1+x)|F'|^2 + \left(\frac{mZ\alpha}{p} \right)^2 \frac{|F|^2}{1-x} \right\}, \\
F &= -\frac{i}{2} \sum_{l=1}^{\infty} l e^{i\pi(l-\nu)} \frac{\Gamma(\nu - iq)}{\Gamma(\nu + 1 + iq)} [P_l(x) - P_{l-1}(x)] \quad , \quad x = \cos \vartheta. \tag{4}
\end{aligned}$$

that coincides with the Mott result [1].

III. LOW-ENERGY SCATTERING

Let us consider now the cross section (4) in the limit $q = Z\alpha\varepsilon/p \gg 1$, $Z\alpha \sim 1$ corresponding to the low-energy electron scattering in the strong Coulomb field. In the first Born

approximation the cross section reads [1]:

$$\frac{d\sigma_B}{d\Omega} = \frac{q^2}{p^2(1-x)^2} \left[1 - \frac{p^2}{2\varepsilon^2}(1-x) \right] \quad . \quad (5)$$

Using the asymptotics of the Γ -function in (4) we represent the ratio $S = d\sigma/d\sigma_B$ in the following form:

$$S = 1 + \frac{(1-x)}{q} \text{Im} \left\{ \exp \left[-iq \ln \left(\frac{1-x}{2} \right) \right] \right. \\ \left. \times \sum_{l=1}^{\infty} l(-1)^l [P_l(x) - P_{l-1}(x)] (e^{-2i\pi\nu} - 1) e^{il^2/q} \right\} . \quad (6)$$

If $(1+x) \gg 1/q$ then it is possible to neglect the factor $\exp(il^2/q)$ in (6). We obtain

$$S = 1 + \frac{(1-x)}{q} \text{Re} \left\{ \exp \left[-iq \ln \left(\frac{1-x}{2} \right) \right] \right. \\ \times \left\{ \pi(Z\alpha)^2 \left(\sqrt{\frac{2}{1+x}} - 1 \right) - i\pi^2(Z\alpha)^4 \ln \sqrt{\frac{1+x}{2}} \right. \\ \left. \left. - i \sum_{l=1}^{\infty} l(-1)^l [P_l(x) - P_{l-1}(x)] \left[e^{-2i\pi\nu} - 1 - \frac{i\pi(Z\alpha)^2}{l} + \frac{\pi^2(Z\alpha)^4}{2l^2} \right] \right\} \right\} \quad . \quad (7)$$

Thus, at $(1+x) \gg 1/q$ the correction to the cross section $\propto 1/q$. Note that the sum in (7) converges very rapidly for any x . When $(1+x) \sim 1/q$ (backward scattering), then the main contribution to the sum in (6) comes from $l \sim \sqrt{q} \gg 1$. Using the asymptotics of the Legendre polynomials at $x \rightarrow -1$ and replacing the summation by the integration we get

$$S = 1 + (1-x) \frac{\pi^{3/2}(Z\alpha)^2}{\sqrt{q}} \cos \left[\frac{\pi}{4} + \frac{q(1+x)}{4} \right] J_0 \left(\frac{q(1+x)}{4} \right) \quad . \quad (8)$$

We see that at $(1+x) \sim 1/q$ the correction to the cross section $\propto 1/\sqrt{q}$. Therefore, for the low-energy scattering the biggest difference between the exact in $Z\alpha$ cross section and $d\sigma_B/d\Omega$ is in the case of backward scattering. For arbitrary q and $x = -1$ the function S reads (see (4)):

$$S = 4 \left| \sum_{l=1}^{\infty} l e^{-i\pi\nu} \frac{\Gamma(\nu - iq)}{\Gamma(\nu + 1 + iq)} \right|^2 \quad . \quad (9)$$

In Fig. 1 we show the dependence of S on $v = p/\varepsilon$ for $x = -1$ and various values of Z , and compare it with the low-energy asymptotics $S = 1 + \sqrt{2} \pi^{3/2} (Z\alpha)^{3/2} \sqrt{v}$.




FIG. 1: The dependence of S on v for $x = -1$ and $Z\alpha = 0.6$ (1), 0.7 (2), and 0.8 (3). Solid lines are the exact results, the dashed lines are the asymptotics.

We see that for backward scattering the exact in $Z\alpha$ cross section differs essentially from $d\sigma_B/d\Omega$ up to rather small v , and this difference decreases very slowly ($\propto \sqrt{v}$). Strictly speaking, the asymptotics are valid when $(S-1) \ll 1$. Nevertheless, it is seen from Fig.1 that the difference between the exact result and the asymptotics is small starting from $v < 0.2$. For $(1+x) \gg 1/q$ one can check that $(S-1) \ll 1$ starting from $q > 10$.

This work was supported through Grant RFBR 01-02-16926.

-
- [1] N.F.Mott, Proc. Roy. Soc. A **124**, 426 (1929).
 - [2] H. Überall, *Electron Scattering From Complex Nuclei* (Academic Press, New York, 1971).
 - [3] J.H. Bartlett, R.E. Watson, Proc. Am. Acad. Arts Sci **74**, 53 (1940).
 - [4] J.A. Doggett, L.V. Spencer, Phys. Rev. **103**, 1597 (1956).
 - [5] N.Sherman, Phys. Rev. **103**, 1601 (1956).
 - [6] A.I. Milstein, V.M. Strakhovenko, Phys. Lett. A **90**, 447 (1982).